

Use of Emergency Medical Services and Timely Treatment Among Ischemic Stroke

Findings From the China Stroke Center Alliance

Hong-Qiu Gu, PhD*; Zhen-Zhen Rao, MPH*; Xin Yang, MD; Chun-Juan Wang, MD; Xing-Quan Zhao, MD; Yi-Long Wang, MD; Li-Ping Liu, MD; Cai-Yun Wang, MD; Chelsea Liu, BA; Hao Li, MD, PhD; Zi-Xiao Li, MD; Rui-Ping Xiao, MD; Yong-Jun Wang, MD; on behalf of the Chinese Stroke Center Alliance Investigators

Background and Purpose—Emergency medical services (EMSs) are critical for early treatment of patients with ischemic stroke, yet data on EMS utilization and its association with timely treatment in China are still limited.

Methods—We examined data from the Chinese Stroke Center Alliance for patients with ischemic stroke from June 2015 to June 2018. Absolute standardized difference was used for covariates' balance assessments. We used multivariable logistic models with the generalized estimating equations to account for intrahospital clustering in identifying demographic and clinical factors associated with EMS use as well as in evaluating the association of EMS use with timely treatment.

Results—Of the 560 447 patients with ischemic stroke analyzed, only 69 841 (12.5%) were transported by EMS. Multivariable-adjusted results indicated that those with younger age, lower levels of education, less insurance coverage, lower income, lower stroke severity, hypertension, diabetes mellitus, and peripheral vascular disease were less likely to use EMS. However, a history of cardiovascular diseases was associated with increased EMS usage. Compared with self-transport, EMS transport was associated with significantly shorter onset-to-door time, door-to-needle time (if prenotification was sent), earlier arrival (adjusted odds ratio [95% CIs] were 2.07 [1.95–2.20] for onset-to-door time ≤ 2 hours, 2.32 [2.18–2.47] for onset-to-door time ≤ 3.5 hours), and more rapid treatment (2.96 [2.88–3.05] for IV-tPA [intravenous recombinant tissue-type plasminogen activator] in eligible patients, 1.70 [1.62–1.77] for treatment with IV-tPA by 3 hours if onset-to-door time ≤ 2 hours, and 1.76 [1.70–1.83] for treatment with IV-tPA by 4.5 hours if onset-to-door time ≤ 3.5 hours).

Conclusions—Although EMS transportation is associated with substantial reductions in prehospital delay and improved likelihood of early arrival and timely treatment, rate of utilization is currently low among Chinese patients with ischemic stroke. Developing an efficient EMS system and promoting culture-adapted education efforts are necessary for improving EMS activation. (*Stroke*. 2019;50:1013-1016. DOI: 10.1161/STROKEAHA.118.024232.)

Key Words: China ■ emergency medical services ■ prehospital delay ■ stroke ■ thrombolytic therapy

Stroke is the leading cause of death in China.¹ Over 70% of prevalent strokes in China are ischemic,² for which thrombolytic therapy is the most effective evidence-based treatment.³ However, prehospital or in-hospital delay may preclude the use of thrombolytics and weaken the treatment effect. Previous studies in high-income countries indicated that emergency medical service (EMS) utilization was independently associated with shorter prehospital delay and more

rapid treatment.⁴⁻⁶ Although several studies reported on EMS utilization in China, they were limited to 1 province or several cities and lacked timely treatment measures.^{7,8} A national program aimed at bridging gaps between guideline recommendations and clinical practices named the Chinese Stroke Center Alliance was sponsored and launched in 2015.⁹ In this study, we address the urgent need to use this national-level data to describe the pattern of EMS utilization, identify associated

Received October 10, 2018; final revision received January 30, 2019; accepted February 12, 2019.

From the China National Clinical Research Center for Neurological Diseases (H.-Q.G., X.Y., C.-J.W., Y.-L.W., C.-Y.W., H.L., Z.-X.L., Y.-J.W.), Vascular Neurology, Department of Neurology (C.-J.W., X.-Q.Z., Y.-L.W., Z.-X.L., Y.-J.W.), Neuro-Intensive Care Unit, Department of Neurology (L.-P.L.), and National Center for Healthcare Quality Management in Neurological Diseases (H.-Q.G., X.Y., C.-J.W., C.-Y.W., Z.-X.L., Y.-J.W.), Beijing Tiantan Hospital, Capital Medical University, China; Institute of Molecular Medicine, Yingjie Center, Peking University, Beijing, China (Z.-Z.R., R.-P.X.); Beijing Key Laboratory of Translational Medicine for Cerebrovascular Disease, China (C.-J.W., Y.-L.W., Y.-J.W.); Center for Stroke, Beijing Institute for Brain Disorders, China (C.-J.W., X.-Q.Z., Y.-L.W., Z.-X.L., Y.-J.W.); and Johns Hopkins Bloomberg School of Public Health, Johns Hopkins University, Baltimore, MD (C.L.).

*Dr Gu and Z.-Z. Rao contributed equally.

The online-only Data Supplement is available with this article at <https://www.ahajournals.org/doi/suppl/10.1161/STROKEAHA.118.024232>.

Correspondence to Zi-Xiao Li, MD, Department of Neurology, Beijing Tiantan Hospital, Capital Medical University, No. 119 Nansihuanxilu, Fengtai District, Beijing 100070, China, Email lizixiao2008@hotmail.com or Rui-Ping Xiao, MD, Institute of Molecular Medicine, Room 322, Ying-Jie Conference Center, Peking University, Yiheyuan Road 5, Haidian District, Beijing 100871, China, Email xiaor@pku.edu.cn

© 2019 American Heart Association, Inc.

Stroke is available at <https://www.ahajournals.org/journal/str>

DOI: 10.1161/STROKEAHA.118.024232

factors with EMS activation, and assess the association between EMS utilization and timeliness of arrival and treatment.

Methods

The data that support the findings of this study are available from the corresponding author upon request.

Details of Chinese Stroke Center Alliance program have been previously reported elsewhere.⁹ All participating hospitals received research approval to collect data without requiring individual patient informed consent under the common rule or a waiver of authorization and exemption from their institutional review board. Study population (Figure I in the [online-only Data Supplement](#)), study variables, and definitions are described in the Methods in the [online-only Data Supplement](#). Given the extensive data set, comparison using $P < 0.05$ indicates statistical significance but may not have any clinical significance. Therefore, baseline characteristics were compared using absolute standardized differences with absolute standardized difference ≥ 10 considered to be clinically significant.¹⁰ We used generalized estimating equations logistic regression modeling with adjustment for within-hospital clustering in both unadjusted and adjusted models. Sensitivity analyses among cases with documented-only or with mode-imputed National Institutes of Health Stroke Scale scores were conducted to determine whether findings are different from primary results.

Results

Among the 560 447 patients with ischemic stroke from 1955 hospitals (Table I in the [online-only Data Supplement](#)), only

69 841 (12.5%) were EMS transported. No regional discrepancies were found (Figure II in the [online-only Data Supplement](#)). Compared with EMS-transported patients, self-transported patients were younger, had lower monthly income, milder stroke severity, and were less likely to have a history of cardiovascular diseases such as atrial fibrillation/flutter, coronary heart disease/myocardial infarction and heart failure (Table II in the [online-only Data Supplement](#)).

All covariates other than gender and carotid stenosis were statistically significantly associated with EMS usage. Strong predictors of EMS usage were cardiovascular diseases, such as atrial fibrillation/flutter, heart failure, coronary heart disease/myocardial infarction, and dyslipidemia. Lower stroke severity was the most substantial barrier to EMS use. Other factors associated with less EMS activation were shown in Figure 1. Similar findings were obtained in sensitivity analyses (Table III in the [online-only Data Supplement](#)).

EMS-transported patients experienced significantly shorter prehospital delay than self-transported patients. However, we did not observe superiority of EMS transportation in the door-to-needle time measure (Table). Nevertheless, if prehospital notification had been sent, there would be fewer minutes of in-hospital delay (61 [40–91] versus 65 [45–95]; $P = 0.0001$). The proportion of patients with an onset-to-door




Figure 1. Factors associated with emergency medical service (EMS) usage. *Adjusted for age, gender, education, insurance, family monthly income per capita, National Institutes of Health Stroke Scale (NIHSS) at admission, history of coronary heart disease (CHD)/myocardial infarction (MI), atrial fib/flutter, stroke/transient ischemic attack (TIA), heart failure, hypertension, carotid stenosis, diabetes mellitus, dyslipidemia, and peripheral vascular disease (PVD) when it is appropriate. NRCMS indicates new rural cooperative medical scheme; UEMIS, urban employ medical insurance scheme; and URMIS, urban residents medical insurance scheme.

Table. Timeline Comparison Between Self-Transport and EMS Transport

Timeline Measure	Eligible Patients	Median (IQR), min		P
		Self-Transport	EMS Transport	
Onset-to-door time	514 140	1300 (243–3099)	271 (109–1379)	<0.0001
Door-to-needle time*	24 542	62 (40–94)	62 (40–93)	0.6994
Onset-to-needle time*	24 542	171 (127–220)	169 (125–215)	0.0002

EMS indicates emergency medical services; and IQR, interquartile range.

*Nine hundred sixty-nine patients with undocumented needle time were excluded from 25 510 IV-tPA (intravenous recombinant tissue-type plasminogen activator) patients from this analysis.

time ≤2 hours among EMS-transported patients was almost 2-fold that of self-transported patients (30.2% versus 15.1%, $P<0.0001$). The adjusted results indicated that compared with self-transported patients, EMS transportation was associated with a 2.07-fold (95% CI, 1.95–2.20) or 2.32-fold (95% CI, 2.18–2.47) higher likelihood of onset-to-door time ≤2 hours or onset-to-door time ≤3.5 hours, respectively. EMS-transported patients were nearly 3× (adjusted odds ratio, 2.96; 95% CI, 2.88–3.05) more likely to receive IV-tPA (intravenous recombinant tissue-type plasminogen activator). Similar results were found in treatment with IV-tPA within 3 hours among eligible patients with onset-to-door time ≤2 hours (adjusted odds ratio, 1.70; 95% CI, 1.62–1.77) or treatment with IV-tPA within 4.5 hours among eligible patients with onset-to-door time ≤3.5 hours (adjusted odds ratio, 1.76; 95% CI, 1.70–1.83) (Figure 2). Sensitivity

analyses yielded similar results (Tables IV and V in the online-only Data Supplement).

Discussion

EMS is strongly recommended for timely arrival and rapid treatment of patients with stroke in the recent guidelines from the American Heart Association and American Stroke Association.¹¹ However, <13% of patients with ischemic stroke were transported to the hospital by EMS in our study, which was much lower than 59.6% in the DASH II Study (The Second Delay in Accessing Stroke Healthcare) or 63.7% in the Get With The Guidelines-Stroke program.^{4,12}

We identified several factors associated with decreased use of EMS, including lower stroke severity, younger age, and lower educational attainment. However, the underlying reason for these characteristics' association with lower EMS utilization cannot be determined by our data at this time. A possible explanation may be the lack of accessibility to EMS or lack of awareness of the need for urgent stroke treatment. Developing a more efficient EMS system and promoting culture-adapted stroke education programs, such as Stroke 1-2-0, on timely recognition of stroke and awareness of the importance of EMS activation are needed to address these disparities.^{13,14}

Our study has some limitations. This observational study is based on hospitals' voluntary enrollment and does not have an elaborately designed sampling frame. However, the large sample size and the generalized estimating equations model help improve the robustness and generalizability of the study. We do not have data on distance traveled from symptom onset to the hospital so we could not assess the relationship between distance and prehospital delay. However, other studies have shown that EMS transportation is associated with




Figure 2. Association between emergency medical service (EMS) and Timely Arrival and Rapid Treatment. *Adjusted for age, gender, National Institutes of Health Stroke Scale (NIHSS) at admission, history of atrial fib/flutter, stroke or transient ischemic attack, heart failure, carotid stenosis, coronary heart disease or myocardial infarction, hypertension, diabetes mellitus, dyslipidemia, and peripheral vascular disease. †Patients (12289; 8921 by self-transport and 3368 by EMS transport) with contraindications to IV-tPA (intravenous recombinant tissue-type plasminogen activator) within the time window were excluded from this analysis. ‡Patients (6037; 4193 by self-transport and 1844 by EMS transport) with contraindications to IV-tPA within the time window were excluded from this analysis. ¶Patients (9448; 6660 by self-transport and 2788 by EMS transport) with contraindications to IV-tPA within the time window were excluded from this analysis. OR indicates odds ratio.

shorter prehospital delay regardless of geographic distance.⁶ Finally, undocumented National Institutes of Health Stroke Scale scores, symptoms or arrival time data may introduce selection bias. To address this issue, Tables VI and VII in the [online-only Data Supplement](#) comparing characteristics were provided and results showed comparable covariates. Results from the sensitivity analysis also confirmed the robustness of our primary analysis.

Conclusions

Our study showed that only 1 in 8 patients with ischemic stroke was EMS transported to the hospital, a proportion that is much lower than that of high-income countries. EMS usage is associated with shorter prehospital delay and a higher likelihood of timely treatment. To improve EMS activation, more emphasis should be placed on developing an efficient EMS system and promoting culture-adapted stroke education programs.

Acknowledgments

We thank all participating centers in the Chinese Stroke Center Alliance program for their hard work in data collection.

Sources of Funding

This work was supported by grants from National Key R&D Program of China (2017YFC1310901, 2016YFC0901001, 2016YFC0901002), National Natural Science Foundation of China (81600999).

Disclosures

None.

References

1. Yang G, Wang Y, Zeng Y, Gao GF, Liang X, Zhou M, et al. Rapid health transition in China, 1990-2010: findings from the Global Burden of Disease Study 2010. *Lancet*. 2013;381:1987-2015. doi: 10.1016/S0140-6736(13)61097-1
2. Wang W, Jiang B, Sun H, Ru X, Sun D, Wang L, et al; NESS-China Investigators. Prevalence, incidence, and mortality of stroke in China: results from a Nationwide Population-Based Survey of 480 687 adults. *Circulation*. 2017;135:759-771. doi: 10.1161/CIRCULATIONAHA.116.025250
3. Anderson CS, Robinson T, Lindley RI, Arima H, Lavados PM, Lee TH, et al; ENCHANTED Investigators and Coordinators. Low-dose versus standard-dose intravenous alteplase in acute ischemic stroke. *N Engl J Med*. 2016;374:2313-2323. doi: 10.1056/NEJMoa1515510
4. Ekundayo OJ, Saver JL, Fonarow GC, Schwamm LH, Xian Y, Zhao X, et al. Patterns of emergency medical services use and its association with timely stroke treatment: findings from Get With the Guidelines-Stroke. *Circ Cardiovasc Qual Outcomes*. 2013;6:262-269. doi: 10.1161/CIRCOUTCOMES.113.000089
5. Mathews R, Peterson ED, Li S, Roe MT, Glickman SW, Wiviott SD, et al. Use of emergency medical service transport among patients with ST-segment-elevation myocardial infarction: findings from the national cardiovascular data registry acute coronary treatment intervention outcomes network registry-Get With the Guidelines. *Circulation*. 2011;124:154-163.
6. Wein TH, Staub L, Felberg R, Hickenbottom SL, Chan W, Grotta JC, et al. Activation of emergency medical services for acute stroke in a non-urban population: the T.L.L. Temple Foundation Stroke Project. *Stroke*. 2000;31:1925-1928.
7. Jiang B, Ru X, Sun H, Liu H, Sun D, Liu Y, et al. Pre-hospital delay and its associated factors in first-ever stroke registered in communities from three cities in China. *Sci Rep*. 2016;6:29795. doi: 10.1038/srep29795
8. Yin X, Yang T, Gong Y, Zhou Y, Li W, Song X, et al. Determinants of emergency medical services utilization among acute ischemic stroke patients in Hubei province in China. *Stroke*. 2016;47:891-894. doi: 10.1161/STROKEAHA.115.011877
9. Wang Y, Li Z, Wang Y, Zhao X, Liu L, Yang X, et al. Chinese stroke center alliance: a national effort to improve healthcare quality for acute stroke and transient ischaemic attack: rationale, design and preliminary findings. *Stroke Vasc Neurol*. 2018;3:256-262.
10. Austin PC. Using the standardized difference to compare the prevalence of a binary variable between two groups in observational research. *Commun Stat Simul Comput*. 2009;38:1228-1234.
11. Powers WJ, Rabinstein AA, Ackerson T, Adeoye OM, Bambakidis NC, Becker K, et al; American Heart Association Stroke Council. 2018 guidelines for the early management of patients with acute ischemic stroke: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. 2018;49:e46-e110. doi: 10.1161/STR.0000000000000158
12. Schroeder EB, Rosamond WD, Morris DL, Evenson KR, Hinn AR. Determinants of use of emergency medical services in a population with stroke symptoms: the Second Delay in Accessing Stroke Healthcare (DASH II) study. *Stroke*. 2000;31:2591-2596.
13. Acker JE III, Pancioli AM, Crocco TJ, Eckstein MK, Jauch EC, Larrabee H, et al; American Heart Association; American Stroke Association Expert Panel on Emergency Medical Services Systems, Stroke Council. Implementation strategies for emergency medical services within stroke systems of care: a policy statement from the American Heart Association/American Stroke Association Expert Panel on Emergency Medical Services Systems and the Stroke Council. *Stroke*. 2007;38:3097-3115. doi: 10.1161/STROKEAHA.107.186094
14. Zhao J, Liu R. Stroke 1-2-0: a rapid response programme for stroke in China. *Lancet Neurol*. 2017;16:27-28. doi: 10.1016/S1474-4422(16)30283-6